

FOURIER ALGEBRAS ON TOPOLOGICAL FOUNDATION *-SEMIGROUPS

MASSOUD AMINI, ALIREZA MEDGHALCHI

ABSTRACT. We introduce the notion of the Fourier and Fourier-Stieltjes algebra of a topological *-semigroup and show that these are commutative Banach algebras. For a class of foundation semigroups, we show that these are preduals of von Neumann algebras.

1. DEFINITIONS AND NOTATIONS

Let S be a locally compact topological semigroup and $M(S)$ be the Banach algebra of all bounded regular Borel measures μ on S with norm and convolution product

$$\|\mu\| = \int_S d|\mu|, \quad \int_S f(x)d(\mu*v)(x) = \int_S \int_S f(xy)d\mu(x)dv(y), (f \in C_0(S), \mu, v \in M(S)).$$

We consider the mappings L_μ and R_μ of S to $M(S)$ defined by

$$L_\mu(x) = \mu * \delta_x, \quad R_\mu(x) = \delta_x * \mu \quad (x \in S, \mu \in M(S))$$

where δ_x is the point mass at x . Then the **semigroup algebra** $L(S)$ consists of those $\mu \in M(S)$ for which $L_{|\mu|}$ and $R_{|\mu|}$ are continuous with respect to the weak

1991 *Mathematics Subject Classification.* 43A37, 43A20.

Key words and phrases. Fourier algebra, Fourier-Stieltjes algebra, foundation semigroups, semigroup C^* -algebra.

This research was supported by Grant 510-2090 of Shahid Behshti University.

topology of $M(S)$. $L(S)$ is a Banach subalgebra of $M(S)$. S is called **foundation** if $\cup\{\text{supp}(\mu) : \mu \in L(S)\}$ is dense in S .

A **representation** of S is a pair $\{\pi, H_\pi\}$ of a Hilbert space H_π and a semigroup homomorphism $\pi : S \rightarrow B(H_\pi)$. We say that π is **(weakly) continuous** if the mappings $x \mapsto \langle \pi(x)\xi, \eta \rangle$ are continuous on S , for all $\xi, \eta \in H_\pi$, and that π is **bounded** if $\|\pi\| = \sup_{x \in S} \|\pi(x)\| < \infty$. A **topological *-semigroup** is a topological semigroup S with a continuous involution $*$: $S \rightarrow S$. In this case, a representation π is called a ***-representation** if moreover $\pi(x^*) = \pi(x)^*(x \in S)$, where the right hand side is the adjoint operator.

Let S be a topological *-semigroup, $\sum(S)$ be the family of all continuous *-representations π of S with $\|\pi\| \leq 1$, and $\sum_0(S)$ be the irreducible elements of $\sum(S)$. The *-representation $\pi_u = \oplus_{\pi \in \sum_0} \pi$ in the Hilbert space $H_u = \oplus_{\pi \in \sum_0} H_\pi$ is called the **universal representation** of S . This is a continuous *-representation with $\|\pi_u\| \leq 1$.

Now let S be a foundation topological *-semigroup with identity. A *-representation $\{\sigma, H\}$ of $L(S)$ is called **non vanishing** if for every $0 \neq \xi \in H$, there exists $\mu \in L(S)$ with $\sigma(\mu)\xi \neq 0$. let $\sum(L(S))$ be the family of all *-representations of $L(S)$ on a Hilbert space which are non vanishing, then by theorem 2.4 and proof of corollary 2.5 in [12], one has a bijective correspondence between $\sum(S)$ and $\sum(L(S))$ via

$$\langle \tilde{\pi}(\mu)\xi, \eta \rangle = \int_S \langle \pi(x)\xi, \eta \rangle d\mu(x) \quad (\mu \in L(S), \xi, \eta \in H_\pi = H_{\tilde{\pi}}).$$

Moreover π and $\tilde{\pi}$ have the same closed invariant subspaces and so π is irreducible if and only if $\tilde{\pi}$ is so, and

$$\begin{aligned}\pi(x)\tilde{\pi}(\mu) &= \tilde{\pi}(\delta_x * \mu) & (x \in S, \mu \in L(S)). \\ \tilde{\pi}(\mu)\pi(x) &= \tilde{\pi}(\mu * \delta_x)\end{aligned}$$

Next, following [14], we consider the family $\Omega(S)$ of all continuous $*$ -representations ω of S in a W^* -algebra with $\|\omega\| \leq 1$ (i.e. semigroup homomorphisms from S into the unit ball of a W^* -algebra M_ω such that $\omega(x^*) = \omega(x)^*$ for all $x \in S$).

2. SEMIGROUP C^* -ALGEBRA AND BIG SEMIGROUP ALGEBRA

Let S be a topological $*$ -semigroup with identity. Given $\rho \subseteq \Sigma = \Sigma(S)$ and $\mu \in L(S)$, define $\|\mu\|_\rho = \sup\{\|\tilde{\pi}(\mu)\| : \pi \in \rho\}$ and $I_\rho = \{\mu \in L(S) : \|\mu\|_\rho = 0\}$. Then I_ρ is clearly a closed two-sided ideal of $L(S)$ and $\|\mu + I_\rho\| = \|\mu\|_\rho$ defines a C^* -norm on $L(S)/I_\rho$. The completion of this quotient space in this norm is a C^* -algebra which is denoted by $C_\rho^*(S)$. When $\rho = \Sigma$, then C^* -algebra $C^*(S) = C_\Sigma^*(S)$ is called the **(full) semigroup C^* -algebra** of S . If S is foundation and Σ separates the points of S , then $L(S)$ is $*$ -semisimple [12], and so $I_\Sigma = \{0\}$. In this case $L(S)$ is a norm dense subalgebra of $C^*(S)$. In particular each $\tilde{\pi} \in \Sigma(L(S))$ uniquely extended to a $*$ -representation of $C^*(S)$ which we still denote it by $\tilde{\pi}$.

Following [14] one may also consider the C^* -algebra $C_\Omega^*(S)$ generated by the set of operators $\{\pi(x) : x \in S, \pi \in \Omega(S)\}$. If $\omega_\Omega = \oplus_{\omega \in \Omega} \omega$ and $M_\Omega = \sum_{\omega \in \Omega} \oplus M_\omega$, then $C_\Omega^*(S)$ is the span closure of $\omega_\Omega(S)$ in M_Ω . We shall define the Fourier-Stieltjes algebra $B(S)$ naturally as the linear span of all continuous positive definite functions on S and one expects that as in the group case [8], the Fourier-Stieltjes algebra is isometrically isomorphic to the dual space of the semigroup C^* -algebra. But $C_\Omega^*(S)^*$

only contains an isometric copy of $B(S)$ in general (see remark 4.3.(e) in [13]), and $C_\Omega^*(S)^* \simeq B(S)$, when S is discrete. As we would show in the next section, if one replaces $C_\Omega^*(S)$ with $C^*(S)$, this pathology disappears and one gets $C^*(S)^* \simeq B(S)$ for each foundation topological $*$ -semigroup with identity for which $\sum(S)$ separates the points of S (theorem 3.3).

Warning: The C^* -algebra $C^*(S)$ defined above is not the same as the one defined in [5], [6]. The reason is that in the latter, one considers representations instead of $*$ -representation. Doing so, $C^*(S)$ could be non commutative even for abelian discrete semigroups, e.g. $C^*(\mathbb{N})$ turns out to be the Toeplitz algebra [16], where as here $C^*(\mathbb{N})$ (when $\mathbb{N}$ is endowed with trivial involution $n^* = n$) is nothing but the algebra $C(\mathbb{D})$ of continuous functions on the unit disk.

Let $W^*(S)$ be the enveloping W^* -algebra of $C^*(S)$. If $\{\pi_u, H_u\}$ is the universal $*$ -representation of S then $\{\tilde{\pi}_u, H_u\}$ is the universal representation of $C^*(S)$ and $W^*(S) = \tilde{\pi}_u(C^*(S))'' \subseteq B(H_u)$ is isometrically isomorphic to $C^*(S)^{**}$ with Arens product [17]. On the other hand, let $W_\Omega^*(S) = \langle \omega_\Omega(S) \rangle^{-\sigma}$ be the weak operator closure of the linear span of $\omega_\Omega(S)$ in M_Ω . Unlike the case of C^* -algebras, one can easily see that the W^* -algebras $W^*(S)$ and $W_\Omega^*(S)$ coincide in general. Indeed to each $\pi \in \sum(S)$, there corresponds a $*$ -representation $\omega_\pi : S \longrightarrow M_\pi = \langle \pi(S) \rangle^{-\sigma} \subseteq B(H_\pi)$ given by $\omega_\pi(x) = \pi(x)$ ($x \in S$). Conversely, if $\omega \in \Omega(S)$ then one can take the universal representation $\{i_\omega, H_\omega\}$ of M_ω to embed M_ω isometrically in $B(H_\omega)$. Then $\pi_\omega = i_\omega \circ \omega \in \sum(S)$, $M_{\pi_\omega} = \langle \pi(S) \rangle^{-\sigma} = \langle i_\omega(\omega(S)) \rangle^{-\sigma} = M_\omega$, and $\omega_{\pi_\omega}(x) = i_\omega(x) = \omega(x)$ ($x \in S$).

In the next proposition, by an abuse of notation we use $\sum(S)$ to denote the set of operators $\{\pi(s) : \pi \in \sum, s \in S\}$ (rather than the set of representations). Also $\sum(L(S)) = \{\tilde{\pi}(\mu) : \pi \in \sum, \mu \in L(S)\}$ and $\sum(C^*(S))$ is defined similarly.

Proposition 2.1. If S is a foundation topological $*$ -semigroup with identity and $\sum$ separates the points of S , then

$$\sum(S)'' = \sum(L(S))'' = \sum(C^*(S))'' = W^*(S),$$

where the commutators are taken in the corresponding universal representations.

Proof. Since $\sum$ separates the points of S , $L(S)$ is norm dense in $C^*(S)$ and so $\sum(L(S))$ is ultra-weakly dense in $\sum(C^*(S))$, hence $\sum(L(S))'' = \sum(C^*(S))'' = W^*(S)$. Now

$$\langle \tilde{\pi}(\mu)\xi, \eta \rangle = \int_S \langle \pi(x)\xi, \eta \rangle d\mu(x) \quad (\mu \in L(S), \xi, \eta \in H_\pi, \pi \in \sum).$$

Hence if $T \in B(H_u)$ commutes with all operators $\pi(x)$, then it also commutes with all operators $\tilde{\pi}(\mu)$, that is $\sum(S)' \subseteq \sum(L(S))'$. Hence $\sum(S)'' \supseteq \sum(L(S))'' = W^*(S)$. On the other hand, for each $x \in S$, $\tilde{\pi}(\delta_x) = \pi(x)$, so $\sum(S)'' \subseteq \sum(\tilde{L}(S))'' = W^*(S)$ and the equality holds. Note that we have used the same notation $\tilde{\pi}$ for the extension of $\tilde{\pi}$ to $M(S)$. $\square$

Proposition 2.2. Let S be a foundation topological $*$ -semigroup with identity and $\sum(S)$ separates the points of S , then $L(S)$, $M(S)$ and $C^*(S)$ are isometrically embedded in $W^*(S)$.

Proof. Consider the universal representation $\{\pi_u, H_u\}$ of S . Since $\sum(S)$ separates the points of S , this is a faithful representation. Hence by theorem 2.4 of

[12], the corresponding representation $\tilde{\pi}_u$ of $L(S)$ is faithful, which in turn is extended to a faithful representation of $C^*(S)$. Also $M(S)$ is the multiplier algebra of $L(S)$ and $L(S)$ has a bounded approximate identity, so $\tilde{\pi}_u$ is extended to a faithful representation of $M(S)$. The theorem follows now from Proposition 2.1. $\square$

Next we study the C^* -algebras $C_\rho^*(S)$, for any $\rho \subseteq \Sigma$ and foundation semigroup S . Consider $\pi \in \Sigma(S)$, then $\tilde{\pi}$ is extended to a $*$ -representation of $M(S)$ on H_π with

$$\tilde{\pi}(\mu) = \int_S \pi(x) d\mu(x) \quad (\mu \in M(S)).$$

If $\rho \subseteq \Sigma$, then $\|\mu\|_\rho = \sup_{\pi \in \rho} \|\tilde{\pi}(\mu)\|$ defines a seminorm on $M(S)$ and for each $\mu, \nu \in M(S)$,

$$\|\mu * \nu\|_\rho \leq \|\mu\|_\rho \|\nu\|_\rho, \quad \|\mu\|_\rho = \|\mu^*\|_\rho, \quad \|\mu * \mu^*\|_\rho = \|\mu\|_\rho^2, \quad \|\mu\|_\rho \leq \|\mu\|_1.$$

Let $N_\rho = \{\mu \in L(S) : \|\mu\|_\rho = 0\}$, then for $\dot{\mu} = \mu + N_\rho \in L(S)/N_\rho$, $\|\dot{\mu}\|_\rho = \|\mu\|_\rho$ defines a norm on $L(S)/N_\rho$ and is independent of the representative μ of the class $\dot{\mu}$, and the completion of $L(S)/N_\rho$ is the C^* -algebra $C_\rho^*(S)$. The next result and its proof are similar to the corresponding result in [8].

Proposition 2.3. For $\rho \subseteq \Sigma$, let $N'_\rho = \bigcap_{\pi \in \rho} \ker \tilde{\pi} \subseteq C^*(S)$, then $N'_\rho \supseteq N_\rho$ and the map $\mu \mapsto \dot{\mu}$ of $L(S)$ onto $L(S)/N_\rho$ is extended uniquely to an isometric epimorphism: $C^*(S) \longrightarrow C_\rho^*(S)$ with kernel N'_ρ ; i.e. $C_\rho^*(S) \simeq C^*(S)/N'_\rho$.

Proof. Since $\|\dot{\mu}\|_\rho \leq \|\mu\|_\Sigma$, the map $\mu \mapsto \dot{\mu}$ is extended continuously to an $*$ -homomorphism $\pi_\rho : C^*(S) \longrightarrow C_\rho^*(S)$ which decreases the norm. Since the image of π_ρ contains $L(S)/N_\rho$, it is dense in $C_\rho^*(S)$. On the other hand, the range of C^* -algebra homomorphisms are always closed [17], so π_ρ is surjective. If $g \in N'_\rho$, then there is a net $\{\mu_\alpha\} \subseteq L(S)$ such that $\|g - \mu_\alpha\|_\Sigma \longrightarrow 0$. Then

$\|\mu_\alpha\|_\rho = \sup_{\pi \in \rho} \|\tilde{\pi}(\mu_\alpha)\| = \sup_{\pi \in \rho} \|\tilde{\pi}(g - \mu_\alpha)\| \leq \|g - \mu_\alpha\|_\Sigma \longrightarrow 0$, hence $\pi_\rho(g) = \lim_\alpha \pi_\rho(\mu_\alpha) = 0$, i.e. $N'_\rho \subseteq \ker \pi_\rho$. Conversely, if $g \in C^*(S)$ and $\pi_\rho(g) = 0$, then there is a net $\{\mu_\alpha\} \subseteq L(S)$ such that $\|\mu_\alpha\|_\rho = \|\dot{\mu}_\alpha\|_\rho \longrightarrow 0$ and $\|g - \mu_\alpha\|_\Sigma \longrightarrow 0$, hence $\sup_{\pi \in \rho} \|\tilde{\pi}(g)\| = \lim_\alpha (\sup_{\pi \in \rho} \|\tilde{\pi}(g) - \tilde{\pi}(\mu_\alpha)\|) \leq \lim_\alpha \|g - \mu_\alpha\|_\Sigma = 0$, i.e. $\tilde{\pi}(g) = 0$ for all $\pi \in \rho$, that's $g \in N'_\rho$. Therefore $\ker \pi_\rho = N'_\rho$. Finally π_ρ is clearly surjective. $\square$

Remark. $\rho \subseteq \Sigma$ separates the points of S exactly when $N_\rho = 0$. This condition is generally weaker than $N'_\rho = 0$.

3. FOURIER AND FOURIER STIELTJES ALGEBRA

Let S be a unital topological $*$ -semigroup. A complex valued function $u : S \longrightarrow \mathbb{C}$ is called **positive definite** if for all positive integers n and all $\lambda_1, \dots, \lambda_n \in \mathbb{C}$, and $x_1, x_2, \dots, x_n \in S$, we have

$$\sum_{i=1}^n \sum_{j=1}^n \lambda_i \bar{\lambda}_j u(x_i x_j^*) \geq 0.$$

Let $P(S)$ denotes the set of all continuous positive definite functions on S . We denote the linear span of $P(S)$ by $B(S)$ and call it the **Fourier Stieltjes algebra** of S .

Proposition 3.1. Let S be a unital topological $*$ -semigroup and $u : S \longrightarrow \mathbb{C}$ be a bounded function, then the followings are equivalent:

- (i) $u \in P(S)$
- (ii) There is a σ -continuous $*$ -representation $\pi : S \longrightarrow B(H)$ and a cyclic vector $\xi \in H$ such that $u = \langle \pi(\cdot)\xi, \xi \rangle$ and $\|\pi\| \leq 1$.

(iii) There is a W^* -algebra M and a σ -continuous $*$ -representation $\omega : S \longrightarrow M$ into the unit ball of M , and a positive bounded normal linear functional $\psi \in M_*$ such that $u = \psi \circ \omega$ and $\langle \omega(S) \rangle^{-\sigma} = M$.

Proof. The proof of equivalence of (i) and (iii) is contained in [14]. Also (ii) $\implies$ (i) is trivial. If (iii) holds, then by Gelfand-Naimark-Segal-construction, there corresponds to $\psi \in M_{*+}$, a nondegenerate σ -continuous $*$ -representation $\{\pi_\psi, H_\psi, \xi_\psi\}$ of M such that $\psi = \langle \pi_\psi(\cdot)\xi_\psi, \xi_\psi \rangle$, and $\overline{\pi_\psi(M)\xi_\psi} = H_\psi$. Let $\pi = \pi_\psi \circ \omega$, then π is a σ -continuous $*$ -representation of S in H_ψ , and $u = \psi \circ \omega = \langle \pi_\psi \circ \omega(\cdot)\xi_\psi, \xi_\psi \rangle = \langle \pi(\cdot)\xi_\psi, \xi_\psi \rangle$, and $\pi(S)\xi_\psi = \pi_\psi(\omega(S))\xi_\psi$ is σ -total in H_ψ . Finally $\|\pi(x)\| = \|\pi_\psi(\omega(x))\| \leq \|\omega(x)\| \leq 1$, for each $x \in S$, and (ii) follows. $\square$

Now consider the algebra $F(S)$ defined by Lau in [14].

Corollary 3.2. If S is a unital topological $*$ -semigroup, then $B(S) = F(S)$. $\square$

Theorem 3.3. Let S be a foundation topological $*$ -semigroup with identity such that $\sum(S)$ separates the points of S . Then $B(S)$ and $C^*(S)^*$ are isometrically isomorphic as Banach spaces.

Proof. Since $\sum(S)$ separates the points of S , $L(S)$ is $*$ -semisimple [11]. In particular $I_\Sigma = \{0\}$, and so $L(S) \subseteq C^*(S)$ is norm dense. Therefore each continuous $*$ -representation of $L(S)$ uniquely extends to one on $C^*(S)$. Now each $u \in B(S) \subseteq C_b(S)$ acts on $L(S)$ via

$$\langle u, \mu \rangle = \int_S u(x) d\mu(x) \quad (\mu \in L(S)).$$

On the other hand, to each $u \in B(S)$, there corresponds some $\pi \in \Sigma(S)$ and $\xi, \eta \in H_\pi$ such that $u(x) = \langle \pi(x)\xi, \eta \rangle$ (Proposition 3.1), and

$$\begin{aligned} \sup\{|\langle u, \mu \rangle| : \mu \in L(S), \|\mu\|_\Sigma \leq 1\} &= \sup_{\|\mu\|_\Sigma \leq 1} \left| \int_S u(x) d\mu(x) \right| \\ &= \sup_{\|\mu\|_\Sigma \leq 1} \left| \int_S \langle \pi(x)\xi, \eta \rangle d\mu(x) \right| \\ &= \sup_{\|\mu\|_\Sigma \leq 1} |\langle \tilde{\pi}(\mu)\xi, \eta \rangle| \\ &\leq \|\tilde{\pi}\| \|\xi\| \|\eta\| \leq \|\xi\| \cdot \|\eta\|. \end{aligned}$$

Hence u extends uniquely to an element of $C^*(S)^*$. Also by density of $L(S)$ in $C^*(S)$, the norm of u as an element of $B(S)$ is the same as the norm of u as an element of $C^*(S)^*$.

Now let $T \in C^*(S)_+^*$, then by restriction, this gives an element of $L(S)_+^*$, hence by the proof of theorem 2.4 (for $\omega = 1$) of [11], there is $\pi \in \Sigma(S)$ and $\xi \in H_\pi$ such that

$$T(\mu) = \int_S \langle \pi(x)\xi, \xi \rangle d\mu(x) \quad (\mu \in L(S)).$$

Define $u \in C_b(S)$ by $u(x) = \langle \pi(x)\xi, \xi \rangle$, then $u \in P(S)$ and $\langle u, \mu \rangle = \int_S \langle \pi(x)\xi, \xi \rangle d\mu(x) = \langle T, \mu \rangle$ ($\mu \in L(S)$), and so $T = u \in P(S)$. $\square$

Theorem 3.4. If S is a unital foundation topological *-semigroup such that $\Sigma(S)$ separates the points of S , then the following norms coincide on $B(S)$ and $B(S)$ is a commutative Banach algebra under each of these norms and $B(S) \simeq W^*(S)_*$, isometrically isomorphic as Banach spaces:

- (i) $\|u\| = \sup\{|\int_S u(x) d\mu(s)| : \mu \in L(S), \sup_{\pi \in \Sigma} \|\tilde{\pi}(\mu)\| \leq 1\},$
- (ii) $\|u\| = \inf\{\|\psi\| : \psi \in M_* \text{ and } u = \psi \circ \omega, \text{ for some } (\omega, M) \in \Omega(S)\},$

- (iii) $\|u\| = \sup\{|\sum_{n=1}^N \lambda_n u(x_n)| : N \geq 1, \lambda_n \in \mathbb{C}, x_n \in S (1 \leq n \leq N),$
 $\sup_{\pi \in \Sigma} \|\sum_{n=1}^N \lambda_n \pi(x_n)\| \leq 1\},$
 (iv) $\|u\| = \inf\{\|\xi\|\|\eta\| : u = \langle \pi(\cdot)\xi, \eta \rangle, \text{ for some } \{\pi, H_\pi\} \in \Sigma, \xi, \eta \in H_\pi\}.$

Proof. It is shown in [14] that $F(S) \simeq W_\Omega^*(S)_*$, with respect to the norm (ii) and in Proposition 3.3 that $B(S) \simeq W^*(S)_*$, with respect to the norm (i). Now $F(S) = B(S)$ and $W_\Omega^*(S)$ and $W^*(S)$ are isometrically isomorphic as W^* -algebras. Hence, by the uniqueness of the predual, the norms (i) and (ii) are the same. Next, by Proposition 2.1, $W^*(S)$ is the W^* -algebra generated by $\{\tilde{\pi}(\delta_x) : x \in S, \pi \in \Sigma(S)\}$. Hence the linear combinations of the form $\sum_{n=1}^N \lambda_n \delta_{x_n}$ are σ -dense in $W^*(S)$. Hence, by Kaplansky's density theorem [17], norms (i) and (iii) coincide. Finally, if $u = \langle \pi(\cdot)\xi, \eta \rangle$, for some $\pi \in \Sigma(S)$ and $\xi, \eta \in H_\pi$, then

$$\begin{aligned} \|u\| &= \sup_{\|\mu\|_\Sigma \leq 1} \left| \int_S u d\mu \right| = \sup_{\|\mu\|_\Sigma \leq 1} \left| \int_S \langle \pi(x)\xi, \eta \rangle d\mu(x) \right| = \sup_{\|\mu\|_\Sigma \leq 1} \left| \langle \tilde{\pi}(\mu)\xi, \eta \rangle \right| \\ &\leq \sup_{\|\mu\|_\Sigma \leq 1} \|\tilde{\pi}(\mu)\| \|\xi\| \|\eta\| \leq \|\xi\| \|\eta\|. \end{aligned}$$

On the other hand, given $\varepsilon > 0$, there is $(\omega, M) \in \Omega(S)$ and $\psi \in M_*$ such that $u = \psi \circ w$ and $\|\psi\| \leq \|u\| + \varepsilon$. Now, by Gelfand-Naimark-Segal-construction, there is a representation $\{\pi_\psi, H_\psi\}$ of M and vector $\xi, \eta \in H_\psi$ such that $\psi = \langle \pi_\psi(\cdot)\xi, \eta \rangle$ and $\|\psi\| = \|\xi\| \|\eta\|$. Put $\pi = \pi_\psi \circ \omega$, then $\{\pi, H_\psi\} \in \Sigma(S)$ and $u = \psi \circ \omega = \langle \pi_\psi \circ \omega(\cdot)\xi, \eta \rangle = \langle \pi(\cdot)\xi, \eta \rangle$, and $\|u\| \geq \|\psi\| - \varepsilon = \|\xi\| \|\eta\| - \varepsilon$. Hence norms (ii) and (iv) also coincide. Now by theorems 3.2 and 4.1 of [14], $B(S)$ is a Banach algebra under pointwise multiplication and norm (ii) and by Theorem 3.3, $B(S) \simeq W^*(S)_*$, under the norm (i), and we are done. $\square$

Proposition 3.5. Let S be a foundation topological $*$ -semigroup with identity, then the followings are equivalent.

- (i) $\Sigma(S)$ separates the points of S ,
- (ii) $B(S)$ separates the points of S ,
- (iii) $L(S)$ is $*$ -semisimple,
- (iv) $M(S)$ is $*$ -semisimple,

Proof. This follows from our Proposition 3.1, and Lemma 3.1 and Theorem 3.4 of [12]. $\square$

Let S be a topological $*$ -semigroup and $C_c(S)$ be the algebra of all continuous functions on S of compact support. Then the closed subalgebra $\overline{B(S) \cap C_c(S)} \subseteq B(S)$ is denoted by $A(S)$ and is called the **Fourier algebra** of S . Since $B(S) \cap C_c(S)$ is clearly a two-sided ideal of $B(S)$ under pointwise multiplication, we have the following result.

Proposition 3.6. $A(S)$ is a closed two-sided ideal of $B(S)$, in particular $A(S)$ is a commutative Banach algebra under pointwise multiplication and each of the norms in Theorem 3.4. $\square$

We observed that $B(S)$ is the predual of a von Neumann algebra. Now we want to see when this holds for $A(S)$. The following lemma should be a part of the literature, but we give a proof for the sake of completeness.

Lemma 3.7. Assume that W is a von Neumann algebra, B is the predual Banach space of W , $z_0 \in W$ is a central projection, and $A = z_0.B$ (Arens product). Then $A^* \simeq z_0 B^*$ (isometrically isomorphic as Banach spaces).

Proof. It is clear that $A \subseteq B$. Let $j : A \longrightarrow B$ be the corresponding linear inclusion, then $j^* : B^* \longrightarrow A^*$ is a surjective continuous linear map and $A^* \simeq B^* / \ker j^*$. Now given $a \in A$ and $f \in B^*$, there is $b \in B$ such that $a = z_0.b$, and so

$(1 - z_0).a = (1 - z_0).(z_0.b) = ((1 - z_0)z_0).b = 0$, hence

$$\langle j^*((1 - z_0)f), a \rangle = \langle (1 - z_0)f, j(a) \rangle = \langle f, (1 - z_0).a \rangle = 0.$$

Hence $(1 - z_0)B^* \subseteq \ker j^*$. Conversely if $f \in \ker j^*$, then for each $b \in B$,

$$\langle z_0f, b \rangle = \langle f, z_0.b \rangle = \langle f, j(z_0.b) \rangle = \langle j^*(f), z_0.b \rangle = 0$$

hence $z_0f = 0$ in B^* ; so $f = (1 - z_0)f \in (1 - z_0)B^*$. Therefore $\ker j^* = (1 - z_0)B^*$

and so $B^*/\ker j^* \simeq z_0B^*$. $\square$

Lemma 3.8. Let S be a foundation topological $*$ -semigroup with identity such that $\sum(S)$ separates the points of S , then $A(S)$ is uniformly dense in $C_0(S)$.

Proof. Let $E = B(S) \cap C_c(S)$. If there is $x \in S$ such that $u(x) = 0$, for each $u \in B(S)$, then for $\delta_x \in W^*(S)$ (see Proposition 2.2)

$$\langle \delta_x, u \rangle = u(x) = 0 \quad (u \in B(S))$$

i.e. $\delta_x = 0$ in $W^*(S)$, which is a contradiction. Hence there is $u \in B(S)$ such that $u(x) \neq 0$, say $u(x) = 1$. Now let $v \in E$ be such that $\|u - v\| < \frac{1}{2}$, then $|u(x) - v(x)| \leq \|u - v\|_\infty \leq \|u - v\| < \frac{1}{2}$, so $|v(x)| \geq \frac{1}{2}$. Therefore E Vanishes nowhere on S . Next, we show that E separates the points of S . Let $x, y \in S$ and $x \neq y$. Then, by Proposition 3.5, there is $u \in B(S)$ such that $u(x) \neq u(y)$. Choose $v \in E$ such that $v(x) \neq 0$, say again $v(x) = 1$. If $v(y) \neq 1$, we have nothing to

proof. Otherwise, $v(x) = v(y) = 1$ and $(uv)(x) \neq (uv)(y)$, and we are done. Now the result follows from Stone-Weierstrauss theorem. $\square$

Theorem 3.9. Let S be a foundation locally compact topological $*$ -semigroup with identity such that $\sum(S)$ separates the points of S , then the followings are equivalent.

- (i) $A(S) \subseteq C^*(S)^* = B(S)$ is invariant under Arens product (i.e. $u.g, g.u \in A(S)$, for each $u \in A(S)$, $g \in C^*(S)$).
- (ii) $A(S) \subseteq W^*(S)_* = B(S)$ is invariant under Arens product (i.e. $u.T, T.u \in A(S)$, for each $u \in A(S)$, $T \in W^*(S)$).
- (iii) $A(S) = I^\perp$, for some w^* -closed ideal I of $W^*(S)$.
- (iv) $A(S) = z_0 \cdot B(S)$, for some central projection $z_0 \in W^*(S)$.
- (v) $A(S)$ is translation invariant.
- (vi) $A(S)^*$ is a W^* -algebra (indeed a quotient of $W^*(S)$).
- (vii) For each compact subset $K \subseteq S$ and each $x \in S$, both Kx^{-1} and $x^{-1}K$ are compact.

Moreover if S is not compact, then these are equivalent to:

- (viii) The one-point compactification S_∞ of S is a semigroup compactification of S .
- (ix) $C_0(S)$ is translation invariant.
- (x) The C^* -algebra $F = C_0(S) \oplus \mathbb{C}$ is an m -admissible subalgebra of S and S_∞ is an F -compactification of S (see [3]).

Proof. The equivalence of (i), (ii), (iii), and (iv) holds in general [4]. Also equivalence of (vii), (viii), (ix), and (x) for the case of locally compact, noncompact

semigroups is contained in [3]. To complete the proof, it is enough to show that

$(iv) \implies (v) \implies (iii)$, $(iv) \implies (vi) \implies (iii)$, and $(vii) \implies (v)$:

$(iv) \implies (v)$. Given $u \in A(S)$, there is $v \in B(S)$ such that $u = v.z_0$. For each $x \in S$, consider $\delta_x \in M(S) \subseteq W^*(S) = B(S)^*$, then

$${}_xu = u.\delta_x = (v.z_0).\delta_x = v.(\delta_x z_0) = v.(z_0 \delta_x) = (v.\delta_x).z_0 = ({}_xv).z_0 \in A(S),$$

since $B(S)$ is always left translation invariant (Theorem 3.2 of [14]).

$(v) \implies (ii)$. By assumption, $\delta_x.u, u.\delta_x \in A(S)$, for each $u \in A(S)$, $x \in S$. But finite sums of the form $\sum_{i=1}^N c_i \delta_{x_i}$ are w^* -dense in $W^*(S)$ (Proposition 2.1), and Arens product is w^* -continuous, hence we get that $A(S)$ is invariant under Arens product by elements of $W^*(S)$.

$(iv) \implies (vi)$. If $A(S) = z_0.B(S)$, then $A(S)^* \simeq z_0 W^*(S)$ is a quotient of the W^* -algebra $W^*(S)$ (lemma 3.7).

$(vi) \implies (iii)$. If $A(S)^*$ is a quotient of $W^*(S)$, then there is a central projection $z_0 \in W^*(S)$ such that $A(S)^* \simeq z_0 W^*(S)$. Take $I = (1 - z_0)W^*(S)$, then $(I^\perp)^* \simeq W^*(S)/I \simeq z_0 W^*(S) \simeq A(S)^*$, so by the uniqueness of the predual of a W^* -algebra, $A(S) \simeq I^\perp$ as Banach spaces.

$(vii) \implies (v)$. Given $u \in A(S)$, there is a net $\{u_\alpha\} \subseteq B(S) \cap C_c(S)$ such that $u_\alpha \longrightarrow u$ in $B(S)$. By the w^* -continuity of the Arens product, for a given $x \in S$, we have ${}_xu_\alpha = \delta_x.u_\alpha \longrightarrow \delta_x.u = {}_xu$, in the norm of $B(S)$. But $\text{supp}({}_xu_\alpha) = x^{-1}(\text{supp } u_\alpha)$ is compact by assumption, that's ${}_xu_\alpha \in B(S) \cap C_c(S)$ for each α , hence ${}_xu \in A(S)$ as required.

$(v) \implies (vii)$ Let $E = B(S) \cap C_c(S)$, then E is dense in $A(S)$ by definition. Also, by the proof of lemma 3.8, E is dense in $C_0(S)$ with respect to the uniform norm. Now given compact subset $K \subseteq S$, there is $g \in C_0(S)$ such that $g = 1$ on K

(Uryshon's lemma). Take $h \in E$ with $\|h - g\|_\infty < \frac{1}{2}$. Then clearly $|h| \geq \frac{1}{2}$ on K . By assumption, ${}_x h \in A(S)$, and so there is $k \in E$ such that in the norm of $A(S)$ we have $\|k - {}_x h\| < \frac{1}{4}$. But $\|k - {}_x h\|_\infty \leq \|k - {}_x h\| < \frac{1}{4}$ (by Theorem 3.2(b) of [4]), and $|{}_x h| \geq \frac{1}{2}$ on $x^{-1}K$, hence $|k| \geq \frac{1}{4}$ on $x^{-1}K$. In particular $x^{-1}K \subseteq \text{supp}(k)$. Since $x^{-1}K$ is compact, therefore $x^{-1}K$ is compact. $\square$

We call a topological semigroup S **(left) weakly cancellative** if it satisfies (the first condition of (iii)) condition (iii) of the above theorem. Note that if S is a Hausdorff cancellative topological semigroup, then it is weakly cancellative (the injective continuous map $y \mapsto yx$ would be a homeomorphism of the compact set K onto Kx^{-1}).

If S is a locally compact, left weakly cancellative, Hausdorff topological semigroup, then it is well known that S has a right Haar measure if and only if it contains a unique minimal left ideal (which is necessarily closed) [1].

If S is a unital, foundation, left weakly cancellative topological $*$ -semigroup, then by the above proposition, $A(S)^*$ is a W^* -algebra quotient of $W^*(S)$. We denote this W^* -algebra by $VN(S)$ and call it the **semigroup von Neumann algebra** of S .

Corollary 3.9. If S is as above, then $W^*(S) \simeq A(S)^\perp \oplus VN(S)$; as W^* -algebras.

Proof. We have

$$A(S)^\perp \simeq (B(S)/A(S))^* \simeq (B(S)/z_0.B(S))^* \simeq (1 - z_0)B(S)^* \simeq (1 - z_0)W^*(S),$$

and $VN(S) \simeq z_0 W^*(S)$. $\square$

Next we consider the problem that when $B(S)$ is the multiplier algebra of $A(S)$. (This is always true for amenable topological groups.) First some notations: Let

A be a Banach algebra and consider A^{**} with first Arens product. This is also a Banach algebra which contains (a copy of) A . Then

$$M(A) = \{x \in A^{**} : xA \cup Ax \subseteq A\}$$

$$\tilde{M}(A) = \{x \in (A^* \cdot A)^* : xA \cup Ax \subseteq A\}$$

are Banach subalgebras of A^{**} containing A . $M(A)$ is called the multiplier algebra of A . If A has a left bounded approximate identity, then $\tilde{M}(A) = RM(A)$, where R stands for right and $RM(A) = \{x \in A^{**} : Ax \subseteq A\}$. In particular, if A moreover is commutative, then $\tilde{M}(A) = M(A)$.

The proof of the next theorem is a modification of the argument given in [15] for topological groups, using the central projection z_0 of the above proposition.

Theorem 3.10. Let S be a unital, foundation, left weakly cancellative topological $*$ -semigroup such that $\sum(S)$ separates the points of S . Then if $A(S)$ has a bounded approximate identity, we have $B(S) = M(A(S))$.

Proof. Consider the map $T : B(S) \longrightarrow (A(S)^* \cdot A(S))^*$ defined by

$$\langle T(a), F \cdot a \rangle = \langle F, ab \rangle \quad (a \in B(S), b \in A(S), F \in A(S)^*).$$

Then T is clearly a linear isometry, and so one may consider $B(S)$ as a Banach subspace of $(A(S)^* \cdot A(S))^*$, where the action on $A(S)$ is the restriction of the product of $B(S)$. Since $A(S)$ is a two sided (closed) ideal of $B(S)$, we have $B(S) \subseteq \tilde{M}(A(S))$. Conversely, if $h \in \tilde{M}(A(S))$, then h is a linear operator on the closed subspace $A(S)^* \cdot A(S)$ of $A(S)^*$, and so it is extended to a linear operator $\tilde{h}$ (with the same norm) on $A(S)^*$ (by Hahn-Banach theorem) such that $a \cdot h = a \cdot \tilde{h} \in A(S)$, for each $a \in A(S)$. Let $\{e_\alpha\} \subseteq A(S)$ be a bounded approximate identity, then passing to

a subset (if necessary) we may assume that $e_\alpha \xrightarrow{w^*} e$, the identity of $A(S)^{**}$, and that $e_\alpha.\tilde{h} \xrightarrow{w^*} \lambda$, for some $\lambda \in B(S) = C^*(S)^*$. Then, given $g \in C^*(S)$, we have

$$\langle g, \lambda \rangle = \lim_{\alpha} \langle g, e_\alpha.h \rangle = \lim_{\alpha} \langle g, e_\alpha.\tilde{h} \rangle = \langle g, e.\tilde{h} \rangle.$$

But, given $a \in A(S)$, we have

$$\langle g.a, e.\tilde{h} \rangle = \langle g, a(e.\tilde{h}) \rangle = \langle g, (ae).\tilde{h} \rangle = \langle g, a.\tilde{h} \rangle = \langle g, a.h \rangle = \langle g.a, h \rangle,$$

hence $e.\tilde{h} = h$, and so $\langle g, \lambda \rangle = \langle g, h \rangle$, for each $g \in C^*(S)$.

Next, given $g \in C^*(S)$ and $a \in A(S)$, we have $g.a \in C^*(S)$ (note that we are now talking about the Arens triple $B(S)$, $B(S)^*$, and $B(S)^{**}$, where as in 3.8(i) we were talking about the triple $C^*(S)$, $C^*(S)^*$, and $C^*(S)^{**}$). Indeed, this follows from the fact that the product of $B(S)$ is a separately w^* -continuous bilinear map. Hence

$$\langle g, a\lambda \rangle = \langle g.a, \lambda \rangle = \langle g.a, h \rangle = \langle g, ah \rangle \quad (g \in C^*(S), a \in A(S)).$$

But $a\lambda, ah \in A(S)$ (since $A(S)$ is an ideal of $B(S)$ and h is a multiplier of $A(S)$), and $A(S) = z_0.B(S)$, hence $z_0.a\lambda = a\lambda$, and $z_0.ah = ah$, for each $a \in A(S)$. Also each element of $A(S)^*$ is of the form z_0T , for some $T \in W^*(S)$, and one can then find a net $\{g_\alpha\} \subseteq C^*(S)$ such that $g_\alpha \xrightarrow{w^*} T$ (since $W^*(S)$ is the enveloping W^* -algebra of $C^*(S)$) to get $z_0g_\alpha \xrightarrow{w^*} F$ and so

$$\begin{aligned} \langle F, a\lambda \rangle &= \lim_{\alpha} \langle z_0g_\alpha, a\lambda \rangle = \lim_{\alpha} \langle g_\alpha, z_0.a\lambda \rangle = \lim_{\alpha} \langle g_\alpha, a\lambda \rangle \\ &= \lim_{\alpha} \langle g_\alpha, ah \rangle = \lim_{\alpha} \langle g_\alpha, z_0.ah \rangle = \lim_{\alpha} \langle z_0g_\alpha, ah \rangle \\ &= \langle F, ah \rangle. \end{aligned}$$

Therefore $\lambda = h$ as elements of $\tilde{M}(A(S))$. In particular $h \in B(S)$ and so $B(S) = \tilde{M}(A(S)) = RM(A(S)) = M(A(S))$. $\square$

REFERENCES

- [1] L. N. Argabright, **A note on invariant integrals on locally compact semigroups**, Proc. Amer. Math. Soc. 17 (1966) 377-382.
- [2] B. A. Barnes, **Representations of the l^1 -algebra of an inverse semigroup**, Trans. Amer. Math. Soc. 218(1976) 361-396.
- [3] J. F. Berglund, D. H. Junghenn, P. Milnes, **Analysis on semigroups**, Candian Math. Soc. Ser. Mono. Adv. Texts, J. Wiley, NY, 1989.
- [4] F. Bonsall, J. Duncan, **Complete normed algebras**, Springer Verlag, Berlin, 1973.
- [5] J. B. Conway, J. Duncan, A. L. T. Paterson, **Monogenic inverse semigroups and their C^* -algebras**, Proc. Roy. Soc. Edinburgh 98A (1984) 13-24.
- [6] J. Duncan, A. L. T. Paterson, **C^* -algebras of inverse semigroups**, Proc. Edinburgh Math. Soc. 28(1985) 41-58.
- [7] J. Ernest, **A new group algebra for locally compact groups**, Amer. J. Math., 86 (1964) 467-492.
- [8] P. Eymard, **L'algebra de Fourier d'un groupe localement compact**, Bull. Soc. Math. France, 92(1964) 181-236.
- [9] J. M. Fell, **The dual spaces of C^* -algebras**, Trans. Amer. Math. Soc., 94(1960) 365-403.
- [10] R. Godement, **Les fonctions de type positive et la theorie des groupes**, Trans. Amer. Math. Soc. 63(1948) 1-84.
- [11] M. Lashkarizadeh Bami, **Bochner's theorem and the Hausdorff moment theorem on foundation topological semigroups**, Can. J. Math. 37(1985) 785-809.
- [12] M. Lashkarizadeh Bami, **Representations of foundation semigroups and their algebras**, Can. J. Math. 37(1985) 29-47.
- [13] A. T. M. Lau, **Analysis on a class of Banach algebras with applications to harmonic analysis on locally compact groups and semigroups**, Found. Math. 68(1983) 161-175.

- [14] A. T. M. Lau, **The Fourier Stieltjes algebra of a topological semigroup with involution**, Pac. J. Math. 77(1978) 165-181.
- [15] A. T. M. Lau, A. Ulger, **Topological centers of certain dual algebras**, Trans. Amer. Math. Soc. 348(1996) 1191-1212.
- [16] G. J. Murphy, **Crossed Products of C^* -algebras by semigroups of automorphisms**, Proc. London Math. Soc. 68(1994) 423-448.
- [17] M. Takesaki, **Theory of operator algebras**, Vol. I, Springer Verlag, 1979.
- [18] M. E. Walter, **W^* -algebras and nonabelian harmonic analysis**, J. Func. Anal. 11(1972) 17-38.

DEPARTMENT OF MATHEMATICS, SHAHID BEHESHTI UNIVERSITY, EVIN, TEHRAN 19839, IRAN,
M-AMINI@CC.SBU.AC.IR

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SASKATCHEWAN, 106 WIGGINS
ROAD, SASKATOON, SASKATCHEWAN, CANADA S7N 5E6, MAMINI@MATH.USASK.CA

DEPARTMENT OF MATHEMATICS, TEACHER TRAINING UNIVERSITY, TEHRAN, IRAN